

This file has been cleaned of potential threats.

If you confirm that the file is coming from a trusted source, you can send the following SHA-256 hash value to your admin for the original file.

d9ee78998bf9fa3954190982bcb16340f4a073c102900c47cffe87f414a990e4

To view the reconstructed contents, please **SCROLL DOWN** to next page.



Decomposition rates of coarse woody debris in undisturbed Amazonian seasonally flooded and unflooded forests in the Rio Negro-Rio Branco Basin in Roraima, Brazil



Reinaldo Imbrozio Barbosa^{a,b,*}, Carolina Volkmer de Castilho^c, Ricardo de Oliveira Perdiz^d, Gabriel Damasco^e, Rafael Rodrigues^f, Philip Martin Fearnside^{b,g}

^a Department of Environmental Dynamics, National Institute for Research in Amazonia (INPA), Roraima Office (NPRR), Rua Coronel Pinto 315 – Centro, 69301-150 Boa Vista, Roraima, Brazil

^b Brazilian Research Network on Climate Change (RedeClima), Brazil

^c Embrapa Roraima, BR 174, Km 8, 69301-970 Boa Vista, Roraima, Brazil

^d Post-graduate Program in Botany (PPG BOT), National Institute for Research in Amazonia (INPA), Av. André Araújo 2936, 69 067-375 Manaus, Amazonas, Brazil

^e Department of Integrative Biology, University of California, Berkeley (UC), 3040 Valley Life Sciences Building # 3140, 94720-3140 Berkeley, CA, USA

^f Post-graduate Program in Tropical Forest Science (PPG CFT), National Institute for Research in Amazonia (INPA), Av. André Araújo 2936, 69 060-001 Manaus, Amazonas, Brazil

^g Department of Environmental Dynamics, National Institute for Research in Amazonia (INPA), Av. André Araújo 2936, 69 067-375 Manaus, Amazonas, Brazil

ARTICLE INFO

Article history:

Received 25 November 2016

Received in revised form 15 April 2017

Accepted 20 April 2017

Keywords:

Carbon flux

Dead biomass

Roraima

Undisturbed forests

Wood density

Wood fragmentation

ABSTRACT

Estimates of carbon-stock changes in forest ecosystems require information on dead wood decomposition rates. In the Amazon, the lack of data is dramatic due to the small number of studies and the large range of forest types. The aim of this study was to estimate the decomposition rate of coarse woody debris (CWD) in two oligotrophic undisturbed forest formations of the northern Brazilian Amazon: seasonally flooded and unflooded. We analyzed 20 arboreal individuals (11 tree species and 3 palm species) with distinct wood-density categories. The mean annual decomposition rate of all samples independent of forest formation ranged from 0.044 to 0.963 yr⁻¹, considering two observation periods (12 and 24 months). The highest rate (0.732 ± 0.206 [SD] yr⁻¹) was observed for the lowest wood-density class of palms, whereas the lowest rate (0.119 ± 0.101 yr⁻¹) was determined for trees with high wood density. In terms of forest formation, the rates values differ when weighted by the wood-density classes, indicating that unflooded forest (0.181 ± 0.083 [SE] yr⁻¹; mean decay time 11–30 years) has a decomposition rate ~19% higher than the seasonally flooded formations (0.152 ± 0.072 yr⁻¹; 13–37 years). This result reflects the dominance of species with high wood density in seasonally flooded formations. In both formations 95% of the dead wood is expected to disappear within 30–40 years. Based on our results, we conclude that the CWD decomposition in the studied area is slower in forests on nutrient-poor seasonally flooded soils, where structure and species composition result in ~40% of the aboveground biomass being in tree species with high wood density. Thus, it is estimated that CWD in seasonally flooded forest formations has longer residence time and slower carbon release by decomposition (respiration) than in unflooded forests. These results improve our ability to model stocks and fluxes of carbon derived from decomposition of dead wood in undisturbed oligotrophic forests in the Rio Negro-Rio Branco Basin, northern Brazilian Amazon.

© 2017 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Greenhouse gases are emitted from tropical forests not only from deforestation but also from standing forests, both disturbed

and undisturbed. Emissions from standing forest have received much less research attention than deforestation. The December 2015 Paris Accords on climate change (UNFCCC, 2015) make it critical to gain better understanding of emissions from standing forest, including the time path of these emissions. The Paris Accords call for preventing mean global temperature from passing a limit “well below” 2 °C the pre-industrial mean, and to “pursue efforts” to keep mean temperature within 1.5 °C of the pre-industrial level.

* Corresponding author at: Department of Environmental Dynamics, National Institute for Research in Amazonia (INPA), Roraima Office (NPRR), Rua Coronel Pinto 315 – Centro, 69301-150 Boa Vista, Roraima, Brazil.

E-mail addresses: reinaldo@inpa.gov.br, imbrozio@gmail.com (R.I. Barbosa).

This means that the net total of all greenhouse gas fluxes, whether anthropogenic or not, must be reduced within a period of approximately 20 years such that the atmospheric concentrations are held close to their present levels (e.g., Rogelj et al., 2016). This requires faster action and greater attention to the time path of emissions than the previous criterion, which defined the limit in terms of a longterm “stabilization” of greenhouse gas concentrations (UNFCCC, 1992, Article 2).

One of the major ways that standing tropical forests emit greenhouse gases is by the death and decomposition of trees, which may be caused by disturbances such as logging, forest fires, invasion by plants such as lianas and bamboos, or by extreme droughts or floods. These disturbances may be either “directly human induced” (in the terms of the Kyoto Protocol: UNFCCC, 1998) or not, as in extreme events influenced by climate change. Coarse woody debris (CWD, diameter ≥ 10 cm) represents a key forest ecosystem compartment in carbon fluxes from standing forest. The carbon stock in this compartment is also important in quantifying the emissions of deforestation, and cannot be simply defined away on the bases of ‘necromass’ not being considered to be ‘biomass’ (e.g., IPCC, 1997); as was done in Brazil’s first and second inventories of greenhouse gases (Brazil-MCT, 2004; Brazil-MCTI, 2010). In the standing forest, the carbon stock in CWD and its decomposition rate will determine the annual carbon budget from this compartment (Harmon et al., 1986).

The CWD decomposition rate (k) is an important biological metric. It is influenced by environmental conditions, wood density and the quality of the soil substrate (Li et al., 2007; Harmon et al., 2011; Russell et al., 2015). Due to slow decomposition and sporadic mortality of trees in forest environments, few studies have addressed the variability in CWD decomposition rates, although this process is a great source of uncertainty in the context of forest dynamics (Harmon et al., 1995; Chambers et al., 2001; Palace et al., 2012). Accurate measurements of CWD decomposition rates are important for determining the magnitude of the carbon stock and the residence time in this forest compartment, providing crucial data for accurate modeling of the effects of climate change on these stocks and on global carbon fluxes (IPCC, 2006; Zell et al., 2009; Bradford et al., 2014).

The paucity of data on CWD decomposition is particularly acute in Amazonia due to the small number of studies, the vastness of the region and the huge range of forest types determined by different environmental conditions and land uses (Chambers et al., 2000; Héroult et al., 2010; Fearnside, 2016). Most studies in undisturbed forests in Amazonia have assumed a steady-state between production and CWD decomposition in order to estimate residence time, stocks and input of carbon through dead tree biomass (Chao et al., 2008; Silva et al., 2016). However, this steady-state has been altered by direct human activity (e.g., selective logging: Keller et al., 2004; Palace et al., 2008), and indirect effects (e.g., increased frequency of surface fires: Alencar et al., 2015; Barni et al., 2015). In both cases, prolonged droughts associated with extreme weather events can maximize the imbalance (Phillips et al., 2009; Vasconcelos et al., 2013).

Selective logging and deforestation provide an increased stock of residual wood pieces and non-commercial trees that are killed as a result of logging (Fearnside, 2000; Aguiar et al., 2016). In this case, the decomposition rates associated with carbon emissions are highly dependent on the replacement landscape (e.g., pastures and secondary forest), with most CWD being considered missing after the first years following deforestation (Buschbacher, 1984; Barbosa and Fearnside, 1996; Fearnside, 2008). In contrast, prolonged drought and surface fires increase tree mortality, reducing standing biomass and dramatically increasing the amount of CWD in undisturbed ecosystems (Brando et al., 2014; Doughty et al., 2015). This imbalance has a direct influence on the residence

time of carbon fixed in the living biomass and its subsequent emission from CWD decomposition in undisturbed forests, and it can become part of a positive feedback process with global climate change (Galbraith et al., 2013). Because the Amazon supports forest formations with high species diversity and large carbon reservoirs (Nogueira et al., 2015), it is crucial to estimate CWD decomposition taking into account specific wood attributes (e.g., wood density) associated with the environmental conditions, and the community structure and species composition of the different forest types (Toledo et al., 2009). This provides a weighted calculation of CWD decomposition rates in terms of forest type, avoiding the use of simple averages that do not represent the ecosystem as a whole.

The Rio Negro–Rio Branco Basin ($\sim 600,000$ km²; Montero and Latrubesse, 2013) lacks information on CWD stocks and carbon flows (Barbosa and Ferreira, 2004; Chao et al., 2009; Silva et al., 2016). This region is characterized by a remarkably variable hydrological regime (different hydro-edaphic restrictions) that shapes vegetation mosaics among seasonally flooded and unflooded forest formations (Junk et al., 2011; Targhetta et al., 2015; Barni et al., 2016). Flooded and unflooded forest types are distinct in community structure and species composition (Hawes et al., 2012), as well as in abiotic characteristics. The present study offers an opportunity to determine the relative importance of taxonomic groups and wood-density classes on CWD decomposition rates in different forests in the Amazon. This approach will improve our understanding of CWD decomposition in undisturbed Amazon forests, especially in ecosystems where distinct hydro-edaphic restrictions can determine differences in the natural decomposition rates.

This study aims to investigate the CWD decomposition rates of different species that occur in undisturbed oligotrophic forests in the Rio Negro–Rio Branco Basin in the Northern Amazon. Our goal was to estimate CWD decomposition rates in two forest formations (seasonally flooded and unflooded) that naturally occur in this region, based on observations made on species in different wood-density classes over two time periods (12 and 24 months). It is expected that CWD decomposition rates in seasonally flooded formations are lower because this type of environment supports tree species with higher wood density and resistance to natural processes of fragmentation (Parolin and Worbes, 2000; Wittmann et al., 2006). The specific objectives of the study were (i) to estimate annual CWD decomposition rates of different taxonomic groups and wood-density classes, (ii) to estimate decomposition rate for each forest formation considering taxonomic groups and wood-density classes weighted by the dominance of individuals (structure) and species (composition) of each formation, and (iii) to estimate the residence time (decay time = number of years for the wood piece to completely lose its physical integrity) and carbon emissions to the atmosphere associated with the annual CWD production in each forest formation. The decomposition rates given here are the first for the oligotrophic forests of the northern Brazilian Amazon.

2. Materials and methods

2.1. Study area

Sampling for CWD decomposition rates was performed in the PPBio (Biodiversity Research Program) research grid in Viruá National Park (1°36'N, 61°13'W); a federal protected area (215,917 ha) located in the state of Roraima (Fig. 1). Viruá is part of the Rio Negro–Rio Branco basin, an ecoregion of the Amazon where the vegetation structure is directly related to the hydro-edaphic restrictions determined by different topographical features, soils and flooding levels (Cordeiro et al., 2016). Viruá is set




Fig. 1. Study area: (A) North of South America; state of Roraima highlighted, (B) Rio Negro-Rio Branco basin (Viruá National Park highlighted) and (C) PPBio grid system installed in Viruá – SRTM image provided by Brazilian Biodiversity Research Program (PPBio, 2014).

in a climatic transition zone (Aw-Am under the Köppen classification system) with a dry season lasting from January to March and a wet season between May and August (Schaefer et al., 2008). Annual rainfall ranges from 1750 to 2000 mm (Barbosa, 1997). In general, the PPBio grid in Viruá can be divided into two major forest formations: (i) seasonally flooded forests characterized by white-sand hydromorphic soils (also called “campinas” and “campinaranas”) and alluvial forests occurring along major watercourses, and (ii) unflooded forests (upland ombrophilous forests) scattered among isolated mountain ranges and ecotone zones, occurring on relatively shallow soils with rocky outcrops (Damasco et al., 2013; Mendonça et al., 2014; Vale Jr. et al., 2016).

2.2. Sampling design

CWD samples to estimate annual decomposition rates were selected from dead arboreal individuals (Trees and Palms) detected by monthly inspections carried out on the PPBio trail grid between December 2011 and March 2012 (t_0 = initial time). The samples were selected based on the following criteria: (i) Trees and Palms with known mortality history (<30 days) determined by the falling of other trees (i.e., living individuals that died due to mechanical action of the wind or from other individuals) and (ii) general bole appearance (bark, heartwood and sapwood) with no trace of fragmentation by decomposition. These criteria were used in order to avoid bias in sampling, ensuring the same initial physical condition and that the date of death is known (the tree falling does not necessarily represent its death or the beginning of decomposition).

We sampled 20 individuals (Trees = 15 individuals in 11 species; Palms = 5 individuals in 3 species) (Appendix A, Table A1). All of these samples were collected in unflooded forest. Taxonomic identification was done in the field for species that could be easily recognized; we did not collect voucher specimens of these species. For individuals for which vouchers were collected, we made the taxonomic determinations by comparison with herbarium specimens housed at MIRR, INPA, RB and UFRR (acronyms following *Index Herbariorum* (Thiers, 2016 [continuously updated])). The

families followed the APG-IV (2016) system; collected specimens were deposited in the UFRR herbarium.

The bole of each selected individual was segmented into 25 discs using a chainsaw, taking into account the distance between the approximate DBH position and the first significant branch (tree canopy base). In total, 500 discs were numbered and the biometric attributes of each piece were measured using calipers (disc width and bark thickness) and a measuring tape (circumference of each disc). The central disc of each individual was taken to the laboratory where it served as a control sample for estimating moisture (water content; %) and basic wood density (ratio between dry weight and saturated volume in the field; g cm^{-3}) at the initial time of observation (t_0). The initial dry weight (initial biomass) of each piece was estimated from the relationship among volume, wet weight and dry weight calculated for each control sample. This was done because it was not possible to remove all discs from the field, dry them in the laboratory and return them back to the field. The basic wood density of each control sample served as the basis for associating the sample with the categories adopted by the Brazilian Institute of the Environment and Renewable Natural Resources (IBAMA) (Sternadt, 2001): very-soft wood species ($\leq 0.350 \text{ g cm}^{-3}$), soft ($0.351\text{--}0.500 \text{ g cm}^{-3}$), medium-soft ($0.501\text{--}0.650 \text{ g cm}^{-3}$), medium-hard ($0.651\text{--}0.800 \text{ g cm}^{-3}$), hard ($0.801\text{--}0.950 \text{ g cm}^{-3}$) and very-hard ($\geq 0.951 \text{ g cm}^{-3}$).

All of the discs that were not controls (i.e., 24 discs per individual) were left at the site where the individual had fallen. These discs were divided into two groups: 12 discs were placed on the ground with the broad (cross-sectional) face in contact with the soil (denominated “latitudinal” discs) and the other 12 discs were placed standing on edge (denominated “longitudinal” discs). The positioning of the discs can be seen in the Appendix A (Fig. A1, panel D). This equitable distribution was adopted to represent different positions of CWD pieces in relation to contact with the soil in the same way as occurs in natural environments. After the initial field experiment had been installed (t_0), two observations were made (t_1 and t_2 , after 12 and 24 months, respectively). Each observation was based on a random collection of four discs from the sampled individual (two latitudinal discs and two longitudinal).

The remaining sampling discs were discarded. The collected discs were dried in an oven at 100 ± 2 °C until constant weight (kg) and were not returned to the field.

2.3. Data analysis

2.3.1. Annual decomposition rate

The annual decomposition rate (k ; yr^{-1} ; \pm SD [standard deviation]) of each individual was estimated as a function of the temporal variation between the initial (t_0) and final (t_1 and t_2) biomasses based on the average of the collected discs (latitudinal + longitudinal) in each observation period, following the model of Olson (1963):

$$k = \frac{-\ln\left(\frac{X_{tn}}{X_0}\right)}{t}$$

where k = annual decomposition rate (fraction per year) of CWD for each individual; X_{tn} = residual biomass (kg) based on the average of four pieces sampled at time t_n (two longitudinal + two latitudinal); X_0 = initial biomass (kg) based on the average of four parts at the initial time (t_0); t = time period (in years) between t_0 and t_n .

The estimate of decomposition rate by wood-density class was calculated as a proportional representation of the velocity of fragmentation of each species group (Trees and Palms), and was analyzed by ANOVA followed by Tukey's test ($\alpha = 0.05$). Regression analysis was performed in order to generate the relational model between wood density (independent) and annual decomposition rate (dependent), taking into account the mean values obtained for each individual of the two taxonomic groups in the two sampling periods (t_1 and t_2).

2.3.2. Decomposition rate by forest formation

To calculate the decomposition rate in terms of forest formation (unflooded and seasonally flooded), we assumed that the annual CWD production is a proportional representation of the above-ground live biomass ($\text{DBH} \geq 10$ cm), as suggested by Silva et al. (2016) for the same study area. This was done because the use of a simple average of the decomposition rate does not represent the whole forest formation due to individual intrinsic factors (e.g., wood density, degree of lignification) that are not incorporated proportionately in the final decomposition rate.

We therefore took into account the results of a forest inventory conducted in the Viruá grid by C.V. de Castilho and collaborators (29 plots of 1 ha: 17 seasonally flooded and 12 unflooded) and produced a database where each individual with $\text{DBH} \geq 10$ cm (Trees and Palms) was associated with its basic wood density based on published databases (Chave et al., 2006; Wittmann et al., 2006; Nogueira et al., 2007; Zanne et al., 2009). After completing this step, the biomass of each individual in the Trees group was estimated using the "moist-forest" model (Chave et al., 2005) taking into account the wood density values associated with the species. The biomasses of individuals without taxonomic identification were calculated using the default value of 0.642 g cm^{-3} for basic wood density (Nogueira et al., 2007). Palm biomass (Arecaceae) was calculated using the Goodman model (Goodman et al., 2013). All biomass calculated from the forest inventory was distributed within its respective wood-density class, being distinguished among taxonomic groups (Trees and Palms) and forest formation (seasonally flooded and unflooded). The biomass of unidentified individuals for trees (5–10% of the total) was proportionally distributed across all wood-density classes described by IBAMA (Sternadt, 2001) in order to preserve the total biomass estimated for the forest formation and to maintain the initial balance among the wood-density classes.

The decomposition rate for each forest formation was estimated by weighting the individual rates of each wood-density class, taking into account the proportional tree biomass distribution of each taxonomic group within each forest formation by observation period. For calculation purposes, the decomposition rate of the Hard and Very-hard classes in the Trees group was estimated by regression based on the values of the other categories. For comparison between the two forest formations, we assume that the decomposition rate obtained in each wood-density class in unflooded forest holds for both forest formations, and the formations were distinguished by their respective proportions of taxonomic groups and wood-density categories. We also assumed no effect in relation to diameter of the sample pieces because the mean diameter of the samples (22.4 ± 6.7 cm; Appendix A, Table A1) was an approximate representation of the mean diameter in the forest inventory (18.2 ± 9.7 cm) conducted in the Viruá grid by C.V. de Castilho and collaborators.

2.3.3. Decay time and CWD carbon flux

Estimates of CWD decay time (\pm SE [standard error]) were based on decomposition rates calculated for each forest formation. For this purpose, the hypothetical CWD decay curve was adopted taking into account the time (independent variable) and the constant of decomposition to 5%, 10%, 25%, 50%, 75% and 95% of the material remaining (dependent variable) as indicated by Olson (1963). Regression analysis was adopted in the hypothetical models for CWD decay time of the two forest formations using a 95% confidence interval (CI). To estimate the long-term annual carbon emissions to the atmosphere (respiration) derived from CWD decomposition, we assumed a predictor interval (range of 95%) of 65–88% of all carbon produced by annual CWD decomposition as indicated by Chambers et al. (2001). The calculations were based on the annual CWD carbon production range (seasonally flooded = $0.04\text{--}0.27 \text{ MgC ha}^{-1} \text{ yr}^{-1}$; unflooded = $0.49\text{--}0.58 \text{ MgC ha}^{-1} \text{ yr}^{-1}$) estimated by Silva et al. (2016) for the PPBio grid in Viruá. All graphics were performed with R software (R Core Team, 2016).

3. Results

3.1. Annual decomposition rate

The annual decomposition rate for all samples varied between 0.044 and 0.963 yr^{-1} , considering the individual average of the latitudinal and longitudinal pieces weighted by the two observation periods (12 and 24 months) (Appendix A, Table A2). On average, decomposition rate of the latitudinal pieces was -18.7% (t_1) and -4.6% (t_2) lower than the rate of the longitudinal pieces. The effect of wood density on the decomposition rates was significant ($\text{ANOVA}_{0.05}$, $F = 15.01$, $p < 0.00001$), indicating that very-soft woods (in low density classes) have higher decomposition rates (Palms = 0.732 ± 0.206 [SD] yr^{-1} ; Trees = $0.630 \pm 0.263 \text{ yr}^{-1}$) as compared to the rates for the higher-density classes (Palms = $0.158 \pm 0.083 \text{ yr}^{-1}$; Trees = $0.119 \pm 0.101 \text{ yr}^{-1}$) (Table 1). The correlation between wood density and decomposition rate in both periods (t_1 and t_2) was negative and significant ($r_p = -0.567$; $p < 0.0001$), and it is best explained by a simple exponential model, especially for the Trees group (Fig. 2).

3.2. Decomposition rate by forest formation

CWD decomposition rate was estimated at 0.152 ± 0.072 [SE] yr^{-1} (CI = $0.081\text{--}0.224 \text{ yr}^{-1}$) for seasonally flooded forests and $0.181 \pm 0.083 \text{ yr}^{-1}$ (CI = $0.098\text{--}0.264 \text{ yr}^{-1}$) for unflooded habitat (Table 2). These results express the structural and floristic differ-

Table 1

Decomposition rate (\pm SD) defined by sampling periods, taxonomic group, wood-density classes and orientation of contact with the soil (CWD, diameter ≥ 10 cm). Different uppercase letters (columns) indicate significant differences between the means of wood-density classes by taxonomic group (ANOVA, Tukey, $\alpha = 0.05$).

Taxonomic group	Wood-density class ^a	n (t ₁ + t ₂)	Latitudinal		Longitudinal		Mean		Weighted mean ^b
			t ₁	t ₂	t ₁	t ₂	t ₁	t ₂	
Trees	Very-soft	8	0.612	0.643	0.630	0.627	0.624 \pm 0.274	0.632 \pm 0.271	0.630 \pm 0.263 ^A
	Soft	6	0.202	0.375	0.219	0.412	0.213 \pm 0.091	0.400 \pm 0.169	0.338 \pm 0.161 ^B
	Medium-soft	8	0.087	0.237	0.238	0.314	0.188 \pm 0.160	0.289 \pm 0.083	0.256 \pm 0.136 ^B
	Medium-hard	8	0.109	0.141	0.120	0.110	0.116 \pm 0.091	0.121 \pm 0.117	0.119 \pm 0.101 ^B
Palms	Very-soft	2	0.893	0.621	1.024	0.606	0.981 \pm 0.093	0.611 \pm 0.010	0.732 \pm 0.206 ^A
	Soft	4	0.225	0.473	0.314	0.439	0.284 \pm 0.126	0.450 \pm 0.056	0.395 \pm 0.135 ^B
	Medium-soft	4	0.070	0.114	0.138	0.211	0.115 \pm 0.043	0.179 \pm 0.109	0.158 \pm 0.083 ^B

^a The decomposition rates (\pm SD) of the Hard ($0.076 \pm 0.073 \text{ yr}^{-1}$) and Very-hard ($0.045 \pm 0.054 \text{ yr}^{-1}$) classes of the Trees group were estimated by regression based on the values of other categories: $Y = 2.141 \times \text{EXP}(-3.514 \times X)$, $R^2 = 0.972$, where Y = annual decomposition rate estimated and X is the upper limit of the each wood-density class described by IBAMA (Sternadt, 2001) (from $X_{\text{very-soft}} = 0.350 \text{ g cm}^{-3}$ to $X_{\text{very-hard}} = 1.100 \text{ g cm}^{-3}$).

^b Considering the average number of days in each period of time (t₁ = 365, t₂ = 743) for all pieces sampled in the field.




Fig. 2. Relation between wood density and CWD decomposition rate (simple exponential model) in each period of time (t₁ and t₂) considering the two taxonomic groups (Palms and Trees) sampled in Viruá National Park, Roraima, Brazil.

ences for formations with different hydro-edaphic restrictions in Viruá, where $\sim 40\%$ of the biomass forming the CWD of the seasonally flooded formation is derived from the higher wood-density classes (Very-hard and Hard) (Appendix A, Tables A3 and A4).

3.3. Decay time and CWD carbon flux

Based on the decomposition rate calculated for each forest formation we estimated that the hypothetical mean time for decomposition of 95% of the CWD produced per year in Viruá is ~ 20 years (CI = 13–37) for the seasonally flooded formation and ~ 17 years (CI = 11–30) for the unflooded formation (Fig. 3). The annual carbon flow to the atmosphere was estimated at $0.002\text{--}0.010 \text{ MgC ha}^{-1} \text{ yr}^{-1}$ (flooded) and $0.022\text{--}0.026 \text{ MgC ha}^{-1} \text{ yr}^{-1}$ (unflooded) considering the CWD carbon production and the respective decay times of the two forest formations. In this scenario, the maximum residence time (upper limit) of estimated CWD carbon is just under 40 years, when it is estimated that 95% of the total necromass will disappear in both formations.

4. Discussion

Our study indicates a considerable variation in annual decomposition rates (range $0.044\text{--}0.963 \text{ yr}^{-1}$) in Viruá. The rates varied according to the taxonomic group and the wood-density class, indicating that the diversity of tree species with different degrees of resistance has a direct impact on the decomposition rates in the study region. The interspecific variation found in Viruá is common in tropical forests, but differs from values found in temperate forests (e.g. Russell et al., 2014; Herrmann et al., 2015) or boreal forests (e.g. Krankina and Harmon, 1995; Freschet et al., 2012), which are ecosystems with lower diversity of species associated with low temperatures and lower CWD decomposition rates.

The high variability in the decomposition rates determined in Viruá was also observed by Martius (1997) in estimates made from wood pieces exposed in a lowland floodplain ecosystem in the central Amazon (range $0.049\text{--}1.000 \text{ yr}^{-1}$). Both cases differ from the higher values found by Chambers et al. (2000) ($0.015\text{--}0.670 \text{ yr}^{-1}$) obtained from estimates derived from tree mortality monitoring in an undisturbed *terra firme* (unflooded upland) forest area in the central Amazon. This difference in upper values is expected for forest formations with different restrictions on the CWD stock (Eaton and Lawrence, 2006). However, distinctions between methods for obtaining the decomposition coefficients also suggest that methodology influences these differences. For example, although there is no firm consensus on the effects of exposure to soil in decomposition rates (e.g. van Geffen et al., 2010), the use of sample pieces in direct contact with the soil in our study influenced the range of values due to increased exposure and the easier access of decomposers to the samples. On the other hand, use of observations derived from monitoring can further reduce the range because most of the samples are derived from large standing dead trees where decomposition rates are initially lower (Palace et al., 2007; Palace et al., 2008).

Regardless of the methods, the most visible implication of our study is the recognition that wood classes with low resistance (Very-soft; $k > 0.600 \text{ yr}^{-1}$) represent trees with rapid decomposition (< 10 years), contrasting with groups with higher resistance ($k < 0.150 \text{ yr}^{-1}$), which contain species with longterm decay attributes. This inverse relationship between decomposition rate and wood density in Viruá was supported by a simple exponential model, as was also observed by Summers (1998) and Chambers et al. (2000) in the central Amazon. While this relationship may be the subject of controversy due to distinct patterns of decomposition among species (Harmon et al., 2000), exponential models have been consistent for Amazonian forests, indicating that they are significant predictors depending on the observed numbers of species and individuals.

Table 2
Decomposition rate (IC; $\alpha = 0.05$) weighted by the average composition of arboreal biomass (diameter ≥ 10 cm; Trees + Palms) of seasonally flooded and unflooded forest formations in Viruá National Park, Roraima, Brazil.

Taxonomic group	Wood-density Class	Flooded		Unflooded	
		Mg ha ⁻¹	k (yr ⁻¹)	Mg ha ⁻¹	k (yr ⁻¹)
Trees	Very-soft	0.29	0.150 (0.079–0.222)	0.57	0.171 (0.092–0.250)
	Soft	9.41		22.08	
	Medium-soft	29.17		40.69	
	Medium-hard	38.26		81.17	
	Hard	48.61		32.20	
	Very-hard	1.84		0.97	
	Sub-total	127.58		177.68	
Palms	Very-soft	0.00	0.195 (0.106–0.284)	3.24	0.386 (0.226–0.546)
	Soft	0.81		0.61	
	Medium-soft	4.36		4.94	
	Medium-hard	0.00		0.00	
	Hard	0.00		0.00	
	Very-hard	0.00		0.00	
	Sub-total	5.16		8.79	
Total	Very-soft	0.29	0.152 (0.081–0.225)	3.81	0.181 (0.098–0.264)
	Soft	10.22		22.69	
	Medium-soft	33.53		45.63	
	Medium-hard	38.26		81.17	
	Hard	48.61		32.20	
	Very-hard	1.84		0.97	
	Total	132.74		186.47	




Fig. 3. Hypothetical time for decomposition of 95% of CWD produced in a year in (A) flooded and (B) unflooded forest formations in Viruá National Park, Roraima, Brazil; CI = confidence interval.

Taxonomic and structural differences between the two forest types (Table 2) are the likely explanation of the $\sim 19\%$ lower decomposition rate for seasonally flooded forests (0.152 yr^{-1}) as compared to that estimated for unflooded forests (0.181 yr^{-1}). Because CWD composition (necromass quality) depends on the forest type and its respective live biomass (Schlegel and Donoso, 2008; Silva et al., 2016), different decomposition rates are expected in distinct regions of the Amazon (Baker et al., 2007), and decomposition rate was best explained in Viruá by the lower abundance of species with high wood density in unflooded forests. The largest weight in the construction of the coefficients was credited to the Trees group (which has greater representation), although the Palms group has higher decomposition rates. The two taxonomic

groups differ in their form of fragmentation, with the center of the Palm samples decomposing more quickly than the outer parts, while the sapwood was lost faster than the heartwood in Trees, in a process similar to that observed by Chao et al. (2008). In any case, the values estimated here were representative of the two formations because the estimates incorporate proportionally the decomposition rates of the different wood-density classes and taxonomic groups. This proportional association generates a proxy for annual CWD production and allows construction of a representative metric and a realistic decay time in order to estimate the carbon flow derived from CWD in each forest formation.

Our estimates for these forests are compatible with the values obtained by Chambers et al. (2000) (0.190 yr^{-1} ; monitoring tree mortality in the central Amazon), and Palace et al. (2008) (0.170 yr^{-1} ; using the steady-state model for large necromass observed in the eastern Amazon). The difference in methods of obtaining decomposition rates indicated no significant effects between the values for the unflooded forests, but there were distinctions in rates for ecosystems with hydro-edaphic restrictions, where forest soils are nutrient-poor (predominantly sandy) and are seasonally flooded.

The highest abundance of high-wood density species partly explain the lower decomposition rate calculated in our study (0.152 yr^{-1}) for seasonally flooded formations, even without considering the effect of the natural deceleration of the decomposition process as a result of seasonal anoxia. Because the decomposition rate of each wood-density class in the flooded forest is based on measurements made in the unflooded forest, these estimates do not include the effect of greatly reduced decomposition rate during the months that the floor of the flooded forest is submerged. If the flooded period were considered, the seasonally flooded forests would tend to be characterized by even lower decomposition rates, as pointed out by Martius (1997), indicating that residence times for CWD carbon in these formations could be even longer than the values estimated here. In this condition, Amazonian seasonally flooded forests can play an important role in the region due their greater proportional capacity for storage of CWD carbon as compared to unflooded forests.

The mean values determined for CWD carbon residence times (17–20 years) in the two formations are within the expected range

for terrestrial forest biomes (12–54 years; Yizhao et al., 2015), but they differ when compared to the studies carried out in the laboratory (e.g. Australian tree species; 6–375 years; Mackensen et al., 2003) or studies on specific groups of trees (e.g. North American hardwoods; 46–71 years; Russell et al., 2014). Although similar to other studies conducted in South American upland rain forests (2–46 years; Hérault et al., 2010), our values are considerably greater than both the default value suggested by the Intergovernmental Panel on Climate Change (IPCC, 1997) (10 years) and the 5–10 years expected for all forest formations in Viruá, when estimated based on the steady-state assumption between stock and annual CWD production (Silva et al., 2016). The assumption of a steady state is commonly used in regional models (Palace et al., 2008), but this method can produce CWD decay times that are unrepresentative when associated with longterm climate change, thereby generating unrealistic estimates for carbon stocks and flows derived from necromass.

Under increasing tree mortality due to forecasted global warming, Amazon forests (~4 milhões km²: Brazil-INPE, 2013) will tend to become in a huge source of carbon emissions to the atmosphere (Brienen et al., 2015; Bonal et al., 2016). In this case, the tendency is for an increase in mortality of larger (non-pioneer) trees that represent substantial amounts of carbon per unit area (Phillips et al., 2010; Lindenmayer et al., 2012), suggesting that undisturbed forests will produce significant amounts of CWD with higher wood density. Within this configuration, the carbon emissions derived from respiration by CWD decomposition in both oligotrophic forest formations in Viruá (flooded = 0.002–0.010 MgC ha⁻¹ yr⁻¹ and unflooded = 0.022–0.026 MgC ha⁻¹ yr⁻¹) will tend to rise due to increasing accumulation of necromass. The magnitude of emissions would be directly related to reducing the carbon sequestration capacity, particularly in seasonally flooded formations due to higher accumulation of CWD with greater resistance to decay as a function of hydro-edaphic characteristics. These differences reflect distinctions in the future CWD carbon stock among the studied formations, directly affecting the disappearance time and the CWD decomposition rate.

This difference in decomposition rates affects the assumptions in Brazil's national inventories of greenhouse gas emissions. The CWD stock and its decomposition are implicitly constant in the third (most recent) inventory (Brazil-MCTI, 2014; Bustamante et al., 2015). However, these stocks are not in equilibrium today, and under a scenario with continued global warming it is expected that tree mortality will increase in undisturbed forests, thereby increasing the stock of CWD and, consequently, emission from decomposition (Barros and Fearnside, 2016). This extra carbon source to the atmosphere is not yet considered by the Brazilian inventory (see Brandão Jr. et al., 2015). Under these conditions, the current net removal rate of carbon estimated for the Amazonia biome (0.43 MgC ha yr⁻¹; Bustamante et al., 2015) will be reduced due to the impoverishment of larger individuals and of species with higher wood density in the forest formations, causing uncertainties in global models and reducing the mitigating role of the region in the global carbon budget.

5. Conclusions

The decomposition rate of coarse woody debris (CWD) in the Viruá area is slower in forests over nutrient-poor and seasonally flooded soils (higher hydro-edaphic restrictions), where the structure and species composition represent ~40% of the aboveground biomass in tree species with high wood density. This means that CWD in seasonally flooded forest formations has a longer residence time and slower carbon release by decomposition (respiration) than in unflooded forests. These results improve our ability to

model carbon stocks and fluxes derived from decomposition of dead wood in undisturbed oligotrophic forests in the Rio Negro-Rio Branco Basin, northern Brazilian Amazon.

Acknowledgments

This study was supported by INPA's institutional project "Ecology and Management of Natural Resources of the Roraima Savanna" (PPI-INPA 015/122). The National Council for Scientific and Technological Development of Brazil (CNPq), provided fellowships for R.I. Barbosa (CNPq 303081/2011-2) and P.M. Fearnside (CNPq 304020/2010-9). The Chico Mendes Institute for Biodiversity Conservation (ICMBio) provided infrastructure and authorization for the study (Authorizations 36441-1 and 36441-2). We also thank CNPq (610042/2009-2, 311103/2015-4) for financial support. This article is a contribution of the Brazilian Research Network on Global Climate Change, FINEP/ Rede CLIMA Grant Number 01.13.0353-00.

Appendix A. Supplementary material

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.foreco.2017.04.026>.

References

- Aguiar, A.P., Vieira, I.C., Assis, T.O., Dalla-Nora, E.L., Toledo, P.M., Santos-Junior, R.A., Batistella, M., Coelho, A.S., Savaget, E.K., Aragão, L.E., Nobre, C.A., Ometto, J.P., 2016. Land use change emission scenarios: anticipating a forest transition process in the Brazilian Amazon. *Global Change Biol.* 22, 1821–1840. <http://dx.doi.org/10.1111/gcb.13134>.
- Alencar, A.A., Brando, P.M., Asner, G.P., Putz, F.E., 2015. Landscape fragmentation, severe drought and the new Amazon forest fire regime. *Ecol. Appl.* 25, 1493–1505. <http://dx.doi.org/10.1890/14-1528.1>.
- APG-IV, 2016. An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG IV. *Bot. J. Linn. Soc.* 181, 1–20. <http://dx.doi.org/10.1111/boj.12385>.
- Baker, T.R., Honorio Coronado, E.N., Phillips, O.L., Martin, J., van der Heijden, G.M., Garcia, M., Silva Espejo, J., 2007. Low stocks of coarse woody debris in a southwest Amazonian forest. *Oecologia* 152, 495–504. <http://dx.doi.org/10.1007/s00442-007-0667-5>.
- Barbosa, R.I., 1997. Distribuição das chuvas em Roraima. In: Barbosa, R.I., Ferreira, E. F.G., Castellon, E.G. (Eds.), *Homem, Ambiente e Ecologia no Estado de Roraima*. Instituto Nacional de Pesquisas da Amazônia (INPA), Manaus, Amazonas, Brazil, pp. 325–335.
- Barbosa, R.I., Fearnside, P.M., 1996. Pasture burning in Amazonia: dynamics of residual biomass and the storage and release of aboveground carbon. *J. Geophys. Res.* 101, 25847–25857. <http://dx.doi.org/10.1029/96JD02090>.
- Barbosa, R.I., Ferreira, C.A.C., 2004. Biomassa acima do solo de um ecossistema de "campina" em Roraima, norte da Amazônia Brasileira. *Acta Amazonica* 34, 577–586. <http://dx.doi.org/10.1590/S0044-59672004000400009>.
- Barni, P.E., Manzi, A.O., Condé, T.M., Barbosa, R.I., Fearnside, P.M., 2016. Spatial distribution of forest biomass in Brazil's state of Roraima, northern Amazonia. *For. Ecol. Manage.* 377, 170–181. <http://dx.doi.org/10.1016/j.foreco.2016.07.010>.
- Barni, P.E., Pereira, V.B., Manzi, A.O., Barbosa, R.I., 2015. Deforestation and forest fires in Roraima and their relationship with phytoclimatic regions in the Northern Brazilian Amazon. *Environ. Manage.* 55, 1124–1138. <http://dx.doi.org/10.1007/s00267-015-0447-7>.
- Barros, H.S., Fearnside, P.M., 2016. Soil carbon stock changes due to edge effects in central Amazon forest fragments. *For. Ecol. Manage.* 379, 30–36. <http://dx.doi.org/10.1016/j.foreco.2016.08.002>.
- Bonal, D., Burban, B., Stahl, C., Wagner, F., Hérault, B., 2016. The response of tropical rainforests to drought—lessons from recent research and future prospects. *Ann. For. Sci.* 73, 27–44. <http://dx.doi.org/10.1007/s13595-015-0522-5>.
- Bradford, M.A., Warren II, R.J., Baldrian, P., Crowther, T.W., Maynard, D.S., Oldfield, E. E., Wieder, W.R., Wood, S.A., King, J.R., 2014. Climate fails to predict wood decomposition at regional scales. *Nat. Climate Change* 4, 625–630. <http://dx.doi.org/10.1038/nclimate2251>.
- Brandão Jr., A., Barreto, P., Souza Jr., C., Brito, B., 2015. Evolução das emissões de gases de efeito estufa no Brasil (1990–2013): Setor de Mudança de Uso da Terra (Documento de Análise). In: SEEG, Observatório do Clima, Instituto do Homem e Meio Ambiente da Amazônia (IMAIZON), Brasília, DF, Brazil. <<http://amazon.org.br/PDFamazon/Portugues/livros/Evolucao%20Emissao%20Gases%20Efeito%20Estufa%20Brasil%201990-2013.pdf>> (accessed 1 October 2015).

- Brando, P.M., Balch, J.K., Nepstad, D.C., Morton, D.C., Putz, F.E., Coe, M.T., Silvério, D., Macedo, M.N., Davidson, E.A., Nóbrega, C.C., Alencar, A., Soares-Filho, B.S., 2014. Abrupt increases in Amazonian tree mortality due to drought–fire interactions. *Proc. Natl. Acad. Sci. USA* <http://dx.doi.org/10.1073/pnas.1305499111>.
- Brazil-INPE, 2013. Projeto PRODES: Monitoramento da Floresta Amazônica Brasileira por Satélite. In: Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, São Paulo, Brazil. Available at: <<http://www.obt.inpe.br/prodes/index.html>> (accessed 25 February 2014).
- Brazil-MCT, 2004. Brazil's Initial National Communication to the United Nations Framework Convention on Climate Change. In: Ministry of Science and Technology (MCT), Brasília, DF, Brazil <<http://unfccc.int/resource/docs/natc/brazilinc1e.pdf>>, p. 271 pp.
- Brazil-MCTI, 2010. Segunda Comunicação Nacional do Brasil à Convenção-Quadro das Nações Unidas sobre Mudança do Clima. In: Coordenação-Geral de Mudanças Globais do Clima, Ministério da Ciência, Tecnologia e Inovação (MCTI), Brasília, DF, Brazil. <http://mct.gov.br/index.php/content/view/326988/Texto_Completo_Publicado.html> (accessed 1 July 2011).
- Brazil-MCTI, 2014. Estimativas anuais de emissões de gases do efeito estufa no Brasil (2a edição). In: Ministério da Ciência, Tecnologia e Inovação, Brasília, DF, Brazil, p. 161. <http://www.mct.gov.br/upd_blob/0235/235580.pdf> (accessed 1 September 2015).
- Brienen, R.J., Phillips, O.L., Feldpausch, T.R., Gloor, E., Baker, T.R., Lloyd, J., Lopez-Gonzalez, G., Monteagudo-Mendoza, A., Malhi, Y., Lewis, S.L., Vasquez Martinez, R., Alexiades, M., Alvarez Davila, E., Alvarez-Loayza, P., Andrade, A., Aragao, L.E., Araujo-Murakami, A., Arets, E.J., Arroyo, L., Aymard, C.G., Banki, O.S., Baraloto, C., Barroso, J., Bonal, D., Boot, R.G., Camargo, J.L., Castilho, C.V., Chama, V., Chao, K.J., Chave, J., Comiskey, J.A., Cornejo Valverde, F., da Costa, L., de Oliveira, E.A., Di Fiore, A., Erwin, T.L., Fauset, S., Forsthofer, M., Galbraith, D.R., Grahame, I.C., Groot, N., Herault, B., Higuchi, N., Honorio Coronado, E.N., Keeling, H., Killeen, T. J., Laurance, W.F., Laurance, S., Licona, J., Magnussen, W.E., Marimon, B.S., Marimon-Junior, B.H., Mendoza, C., Neill, D.A., Nogueira, E.M., Nunez, P., Pallqui Camacho, N.C., Parada, A., Pardo-Molina, G., Peacock, J., Pena-Claros, M., Pickavance, G.C., Pitman, N.C., Poorter, L., Prieto, A., Quesada, C.A., Ramirez, F., Ramirez-Angulo, H., Restrepo, Z., Roosind, A., Rudas, A., Salomao, R.P., Schwarz, M., Silva, N., Silva-Espejo, J.E., Silveira, M., Stropp, J., Talbot, J., ter Steege, H., Teran-Aguilar, J., Terborgh, J., Thomas-Caesar, R., Toledo, M., Tolleiro-Raventos, M., Umetsu, R.K., van der Heijden, G.M., van der Hout, P., Guimarães Vieira, I.C., Vieira, S.A., Vilanova, E., Vos, V.A., Zagt, R.J., 2015. Long-term decline of the Amazon carbon sink. *Nature* 519, 344–348. <http://dx.doi.org/10.1038/nature14283>.
- Buschbacher, R.J., 1984. Changes in Productivity and Nutrient Cycling Following Conversion of Amazon Rainforest to Pasture. University of Georgia, Athens, GA, USA.
- Bustamante, M., Santos, M.M.O., Shimbo, J.Z., Cantinho, R.Z., Bandeira de Mello, T.R., Carvalho e Oliveira, P.V., Cunha, P.W.P., Martins, F.S.R.V., Aguiar, A.P.D., Ometto, J., 2015. Terceiro Inventário Brasileiro de Emissões e Remoções Antrópicas de Gases do Efeito Estufa (Relatórios de Referência): Setor Uso da Terra, Mudança do Uso da Terra e Florestas. In: Ministério da Ciência, Tecnologia e Inovação (MCTI), Brasília, DF, Brazil. <[http://sirene.mcti.gov.br/documents/1686653/1706165/RR_LULUCF_Mudanc%C3%A7%C3%A7%C3%A7%20de+Uso+e+Floresta.pdf/11dc4491-65c1-4895-a8b6-e96705f2717a](http://sirene.mcti.gov.br/documents/1686653/1706165/RR_LULUCF_Mudanc%C3%A7%C3%A7%C3%A7%C3%A7%20de+Uso+e+Floresta.pdf/11dc4491-65c1-4895-a8b6-e96705f2717a)> (accessed 1 December 2015).
- Chambers, J.Q., Higuchi, N., Schimel, J.P., Ferreira, L.V., Melack, J.M., 2000. Decomposition and carbon cycling of dead trees in tropical forests of the Central Amazon. *Oecologia* 122, 380–388. <http://dx.doi.org/10.1007/s004420050044>.
- Chambers, J.Q., Schimell, J.P., Nobre, A.D., 2001. Respiration from coarse wood litter in central Amazon forests. *Biogeochemistry* 52, 115–131. <http://dx.doi.org/10.3334/ORNLAAC/911>.
- Chao, K.-J., Phillips, O.L., Baker, T.R., 2008. Wood density and stocks of coarse woody debris in a northwestern Amazonian landscape. *Can. J. For. Res.* 38, 795–805. <http://dx.doi.org/10.1139/x07-163>.
- Chao, K.-J., Phillips, O.L., Baker, T.R., Peacock, J., Lopez-Gonzalez, G., Martinez, R.V., Monteagudo, A., Torres-Lezama, A., 2009. After trees die: quantities and determinants of necromass across Amazonia. *Biogeosciences* 6, 1615–1626. <http://dx.doi.org/10.5194/bg-6-1615-2009>.
- Chave, J., Andalo, C., Brown, S., Cairns, M.A., Chambers, J.Q., Eamus, D., Fölster, H., Fromard, F., Higuchi, N., Kira, T., Lescure, J.-P., Nelson, B.W., Ogawa, H., Puig, H., Riéra, B., Yamakura, T., 2005. Tree allometry and improved estimation of carbon stocks and balance in tropical forests. *Oecologia* 145, 87–99. <http://dx.doi.org/10.1007/s00442-0050100-x>.
- Chave, J., Muller-Landau, H.C., Baker, T.R., Easdale, T.A., Steege, H.T., Webb, C.O., 2006. Regional and phylogenetic variation of wood density across 2456 Neotropical tree species. *Ecol. Appl.* 16, 2356–2367. [http://dx.doi.org/10.1890/1076-1071\(2006\)016\[2356:RAPVOW\]2.0.CO;2](http://dx.doi.org/10.1890/1076-1071(2006)016[2356:RAPVOW]2.0.CO;2).
- Cordeiro, C.L.O., Rossetti, D.F., Gribel, R., Tuomisto, H., Zani, H., Ferreira, C.A.C., Coelho, L., 2016. Impact of sedimentary processes on white-sand vegetation in an Amazonian megafan. *J. Trop. Ecol.* 32, 498–509. <http://dx.doi.org/10.1017/s0266467416000493>.
- Damasco, G., Vicentini, A., Castilho, C.V., Pimentel, T.P., Nascimento, H.E.M., 2013. Disentangling the role of edaphic variability, flooding regime and topography of Amazonian white-sand vegetation. *J. Veg. Sci.* 24, 384–394. <http://dx.doi.org/10.1111/j.1654-1103.2012.01464.x>.
- Doughty, C.E., Metcalfe, D.B., Girardin, C.A., Amezcua, F.F., Cabrera, D.C., Huasco, W.H., Silva-Espejo, J.E., Araujo-Murakami, A., da Costa, M.G., Rocha, W., Feldpausch, T.R., Mendoza, A.L., da Costa, A.C., Meir, P., Phillips, O.L., Malhi, Y., 2015. Drought impact on forest carbon dynamics and fluxes in Amazonia. *Nature* 519, 78–82. <http://dx.doi.org/10.1038/nature14213>.
- Eaton, J.M., Lawrence, D., 2006. Woody debris stocks and fluxes during succession in a dry tropical forest. *For. Ecol. Manage.* 232, 46–55. <http://dx.doi.org/10.1016/j.foreco.2006.05.038>.
- Fearnside, P.M., 2000. Global warming and tropical land-use change: greenhouse gas emissions from biomass burning, decomposition and soils in forest conversion, shifting cultivation and secondary vegetation. *Climatic Change* 46, 115–158. <http://dx.doi.org/10.1023/A:1005569915357>.
- Fearnside, P.M., 2008. Deforestation in Brazilian Amazonia and global warming. *Ann. Arid Zone* 47, 1–20.
- Fearnside, P.M., 2016. Brazil's Amazonian forest carbon: the key to Southern Amazonia's significance for global climate. *Reg. Environ. Change* <http://dx.doi.org/10.1007/s10113-016-1007-2>.
- Freschet, G.T., Weedon, J.T., Aerts, R., van Hal, J.R., Cornelissen, J.H.C., 2012. Interspecific differences in wood decay rates: insights from a new short-term method to study long-term wood decomposition. *J. Ecol.* 100, 161–170. <http://dx.doi.org/10.1111/j.1365-2745.2011.01896.x>.
- Galbraith, D., Malhi, Y., Affum-Baffoe, K., Castanho, A.D.A., Doughty, C.E., Fisher, R.A., Lewis, S.L., Peh, K.S.-H., Phillips, O.L., Quesada, C.A., Sonké, B., Lloyd, J., 2013. Residence times of woody biomass in tropical forests. *Plant Ecol. Divers.* 6, 139–157. <http://dx.doi.org/10.1080/17550874.2013.770578>.
- Goodman, R.C., Phillips, O.L., del Castillo Torres, D., Freitas, L., Cortese, S.T., Monteagudo, A., Baker, T.R., 2013. Amazon palm biomass and allometry. *For. Ecol. Manage.* 310, 994–1004. <http://dx.doi.org/10.1016/j.foreco.2013.09.045>.
- Harmon, M.E., Franklin, J.F., Swanson, F.J., Sollins, P., Gregory, S.V., Lattin, J.D., Anderson, N.H., Cline, S.P., Aumen, N.G., Sedell, R., Lienkaemper, G.W., Cromack Jr., K., Cummins, K.W., 1986. Ecology of coarse woody debris in temperate ecosystems. *Adv. Ecol. Res.* 15, 133–302. [http://dx.doi.org/10.1016/S0065-2504\(08\)60121-X](http://dx.doi.org/10.1016/S0065-2504(08)60121-X).
- Harmon, M.E., Krankina, O.N., Sexton, J., 2000. Decomposition vectors: a new approach to estimating woody detritus decomposition dynamics. *Can. J. For. Res.* 30, 76–84. <http://dx.doi.org/10.1139/x99-187>.
- Harmon, M.E., Whigham, D.F., Sexton, J., Olmsted, J., 1995. Decomposition and mass of wood detritus in the dry tropical forests of the Northeastern Yucatan Peninsula, Mexico. *Biotropica* 27, 305–316. <http://dx.doi.org/10.2307/2388916>.
- Harmon, M.E., Woodall, C.W., Fasth, B., Sexton, J., Yatkov, M., 2011. Differences between standing and downed dead tree wood density reduction factors: a comparison across decay classes and tree species. In: *Res. Pap. NRS-15*. Department of Agriculture, Forest Service, Northern Research Station, Newtown Square, PA, USA, p. 40 <http://www.nrs.fs.fed.us/pubs/rp/rp_nrs15.pdf>.
- Hawes, J.E., Peres, C.A., Riley, L.B., Hess, L.L., 2012. Landscape-scale variation in structure and biomass of Amazonian seasonally flooded and unflooded forests. *For. Ecol. Manage.* 281, 163–176. <http://dx.doi.org/10.1016/j.foreco.2012.06.023>.
- Héroult, B., Beauchene, J., Muller, F., Wagner, F., Baraloto, C., Blanc, L., Martin, J.M., 2010. Modeling decay rates of dead wood in a neotropical forest. *Oecologia* 164, 243–251. <http://dx.doi.org/10.1007/s00442-010-1602-8>.
- Herrmann, S., Kahl, T., Bauhus, J., 2015. Decomposition dynamics of coarse woody debris of three important central European tree species. *For. Ecosyst. 2*, 27. <http://dx.doi.org/10.1186/s40663-015-0052-5>.
- IPCC, 1997. Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories. In: National Greenhouse Gas Inventories Programme. Intergovernmental Panel on Climate Change (IPCC) and Institute for Global Environmental Strategies (IGES), Kanagawa, Japan <<http://www.ipcc-nggip.iges.or.jp/public/gl/invs6.html>>.
- IPCC, 2006. 2006 IPCC Guidelines for National Greenhouse Gas Inventories. In: Eggleston H.S., B.L., Miwa K., Ngara T., Tanabe K. (Eds.), National Greenhouse Gas Inventories Programme (Agriculture, Forestry and Other Land Use). Intergovernmental Panel on Climate Change (IPCC) and Institute for Global Environmental Strategies (IGES), Kanagawa, Japan. <<http://www.ipcc-nggip.iges.or.jp/public/2006gl/>>.
- Junk, W.J., Piedade, M.T.F., Schöngart, J., Cohn-Haft, M., Adeney, J.M., Wittmann, F., 2011. A classification of major naturally-occurring Amazonian lowland wetlands. *Wetlands* 31, 623–640. <http://dx.doi.org/10.1007/s13157-011-0190-7>.
- Keller, M., Palace, M., Asner, G.P., Pereira, R., Silva, J.N.M., 2004. Coarse woody debris in undisturbed and logged forests in the eastern Brazilian Amazon. *Global Change Biol.* 10, 784–795. <http://dx.doi.org/10.1111/j.1529-8817.2003.00770.x>.
- Krankina, O.N., Harmon, M.E., 1995. Dynamics of the dead wood carbon in Northwestern Russian boreal forests. *Water Air Soil Pollut.* 82, 227–238. <http://dx.doi.org/10.1007/BF01182836>.
- Li, Z., Li-min, D., Hui-yan, G., Lei, Z., 2007. Review on the decomposition and influence factors of coarse woody debris in forest ecosystem. *J. For. Res.* 18, 48–54. <http://dx.doi.org/10.1007/s11676-007-0009-9>.
- Lindenmayer, D.B., Laurance, W.F., Franklin, J.F., 2012. Global decline in large old trees. *Science* 338, 1305–1306. <http://dx.doi.org/10.1126/science.1231070>.
- Mackensen, J., Bauhus, J., Webber, E., 2003. Decomposition rates of coarse woody debris - a review with particular emphasis on Australian tree species. *Aust. J. Bot.* 51, 27–37. <http://dx.doi.org/10.1186/s40663-015-0052-5>.
- Martius, C., 1997. Decomposition of wood. In: Junk, W. (Ed.), *The Central Amazon Floodplain: Ecology of a Pulsing System*. Springer, Heidelberg, Germany, pp. 267–276. http://dx.doi.org/10.1007/978-3-662-03416-3_12.
- Mendonça, B.A.F., Simas, F.N.B., Schaefer, C.E.G.R., Fernandes Filho, E.I., Vale Júnior, J. F., Mendonça, J.G.F., 2014. Podzolized soils and paleoenvironmental

- implications of white-sand vegetation (Campinarana) in the Viruá National Park, Brazil. *Geoderma Reg.* 2–3, 9–20. <http://dx.doi.org/10.1016/j.geodrs.2014.09.004>.
- Montero, J.C., Latrubesse, E.M., 2013. The igapó of the Negro River in central Amazonia: linking late-successional inundation forest with fluvial geomorphology. *J. S. Am. Earth Sci.* 46, 137–149. <http://dx.doi.org/10.1016/j.jsames.2013.05.009>.
- Nogueira, E., Fearnside, P., Nelson, B., Franca, M., 2007. Wood density in forests of Brazil's 'arc of deforestation': implications for biomass and flux of carbon from land-use change in Amazonia. *For. Ecol. Manage.* 248, 119–135. <http://dx.doi.org/10.1016/j.foreco.2007.04.047>.
- Nogueira, E.M., Yanai, A.M., Fonseca, F.O., Fearnside, P.M., 2015. Carbon stock loss from deforestation through 2013 in Brazilian Amazonia. *Global Change Biol.* 21, 1271–1292. <http://dx.doi.org/10.1111/gcb.12798>.
- Olson, J.S., 1963. Energy storage and the balance of producers and decomposers in ecological systems. *Ecology* 44, 322–331. <http://dx.doi.org/10.2307/1932179>.
- Palace, M., Keller, M., Asner, G.P., Silva, J.N.M., Passos, C., 2007. Necromass in undisturbed and logged forests in the Brazilian Amazon. *For. Ecol. Manage.* 238, 309–318. <http://dx.doi.org/10.1016/j.foreco.2006.10.026>.
- Palace, M., Keller, M., Hurr, G., Frolking, S., 2012. A review of above ground necromass in tropical forests. In: Sudarshana, P., Nageswara-Rao, M., Soneji, J.R. (Eds.), *Tropical Forests*. InTech, Rijeka, Croatia, pp. 215–252. <http://dx.doi.org/10.5772/1410>.
- Palace, M., Keller, M., Silva, H., 2008. Necromass production: studies in undisturbed and logged Amazon forests. *Ecol. Appl.* 18, 873–884. <http://dx.doi.org/10.1890/06-2022.1>.
- Parolin, P., Worbes, M., 2000. Wood density of trees in black water floodplains of Rio Jau National Park, Amazonia, Brazil. *Acta Amazonica* 30, 441–448. <http://dx.doi.org/10.1590/1809-43922000303448>.
- Phillips, O.L., Aragão, L.E., Lewis, S.L., Fisher, J.B., Lloyd, J., Lopez-Gonzalez, G., Malhi, Y., Monteagudo, A., Peacock, J., Quesada, C.A., van der Heijden, G., Almeida, S., Amaral, I., Arroyo, L., Aymard, G., Baker, T.R., Banki, O., Blanc, L., Bonal, D., Brando, P., Chave, J., de Oliveira, A.C., Cardozo, N.D., Czimczik, C.I., Feldpausch, T. R., Freitas, M.A., Gloor, E., Higuchi, N., Jimenez, E., Lloyd, G., Meir, P., Mendoza, C., Morel, A., Neill, D.A., Nepstad, D., Patino, S., Penuela, M.C., Prieto, A., Ramirez, F., Schwarz, M., Silva, J., Silveira, M., Thomas, A.S., Steege, H.T., Stropp, J., Vasquez, R., Zelazowski, P., Alvarez Davila, E., Andelman, S., Andrade, A., Chao, K. J., Erwin, T., Di Fiore, A., Honorio, C.E., Keeling, H., Killeen, T.J., Laurance, W.F., Pena Cruz, A., Pitman, N.C., Nunez Vargas, P., Ramirez-Angulo, H., Rudas, A., Salamao, R., Silva, N., Terborgh, J., Torres-Lezama, A., 2009. Drought sensitivity of the Amazon rainforest. *Science* 323, 1344–1347. <http://dx.doi.org/10.1126/science.1164033>.
- Phillips, O.L., van der Heijden, G., Lewis, S.L., López-González, G., Aragão, L.E.O.C., Lloyd, J., Malhi, Y., Monteagudo, A., Almeida, S., Dávila, E.A., Amaral, I., Andelman, S., Andrade, A., Arroyo, L., Aymard, G., Baker, T.R., Blanc, L., Bonal, D., Oliveira, Á.C.A., Chao, K.-J., Cardozo, N.D., Costa, L., Feldpausch, T.R., Fisher, J. B., Fyllas, N.M., Freitas, M.A., Galbraith, D., Gloor, E., Higuchi, N., Honorio, E., Jiménez, E., Keeling, H., Killeen, T.J., Lovett, J.C., Meir, P., Mendoza, C., Morel, A., Vargas, P.N., Patiño, S., Peh, K.S.-H., Cruz, A.P., Prieto, A., Quesada, C.A., Ramirez, F., Ramirez, H., Rudas, A., Salamão, R., Schwarz, M., Silva, J., Silveira, M., Slik, J.W. F., Sonké, B., Thomas, A.S., Stropp, J., Taplin, J.R.D., Vásquez, R., Vilanova, E., 2010. Drought-mortality relationships for tropical forests. *New Phytologist* 187, 631–646. <http://dx.doi.org/10.1111/j.1469-8137.2010.03359.x>.
- PPBio, 2014. Programa de Pesquisa em Biodiversidade: Repositório de dados do PPBio (Mapas - SIG). In: PPBio, CENBAM, Manaus, AM. <<https://ppbio.inpa.gov.br/mapas>>.
- R Core Team, 2016. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria <<http://www.R-project.org/>>.
- Rogelj, J., den Elzen, M., Höhne, N., Fransen, T., Fekete, H., Winkler, H., Schaeffer, R., Fu Sha, R.K., Meinshausen, M., 2016. Paris agreement climate proposals need a boost to keep warming well below 2 °C. *Nature* 534, 631–639. <http://dx.doi.org/10.1038/nature18307>.
- Russell, M.B., Fraver, S., Aakala, T., Gove, J.H., Woodall, C.W., D'Amato, A.W., Ducey, M.J., 2015. Quantifying carbon stores and decomposition in dead wood: a review. *For. Ecol. Manage.* 350, 107–128. <http://dx.doi.org/10.1016/j.foreco.2015.04.033>.
- Russell, M.B., Woodall, C.W., Fraver, S., D'Amato, A.W., Domke, G.M., Skog, K.E., 2014. Residence times and decay rates of downed woody debris biomass/carbon in Eastern US Forests. *Ecosystems* 17, 765–777. <http://dx.doi.org/10.1007/s10021-014-9757-5>.
- Schaefer, C.E.G.R., Mendonça, B.A.F., Fernandes-Filho, E.I., 2008. Geoambientes e paisagens do Parque Nacional do Viruá - RR: Esboço de integração da geomorfologia, climatologia, solos, hidrologia e ecologia (Zoneamento Preliminar). In: Universidade Federal de Viçosa (UFV), Viçosa, Minas Gerais, Brazil, p. 56.
- Schlegel, B.C., Donoso, P.J., 2008. Effects of forest type and stand structure on coarse woody debris in old-growth rainforests in the Valdivian Andes, south-central Chile. *For. Ecol. Manage.* 255, 1906–1914. <http://dx.doi.org/10.1016/j.foreco.2007.12.013>.
- Silva, L.F.S.G., Castilho, C.V., Cavalcante, C.O., Pimentel, T.P., Fearnside, P.M., Barbosa, R.I., 2016. Production and stock of coarse woody debris across a hydro-edaphic gradient of oligotrophic forests in the northern Brazilian Amazon. *For. Ecol. Manage.* 364, 1–9. <http://dx.doi.org/10.1016/j.foreco.2015.12.045>.
- Sternadt, G.H., 2001. *Trabalhabilidade De 108 Espécies De Madeiras Da Região Amazônica*. Instituto Brasileiro do Meio Ambiente e dos Recursos Naturais Renováveis, Laboratório de Produtos Florestais, Brasília, DF.
- Summers, P.M., 1998. Estoque, decomposição e nutrientes da liteira grossa em florestas de terra-firme na Amazônia Central. In: Programa de Pós-graduação em Ciências de Florestas Tropicais. Instituto Nacional de Pesquisas da Amazônia (INPA) e Universidade do Amazonas (UA), Manaus, Amazonas, Brazil, p. 136.
- Targhetta, N., Kesselmeier, J., Wittmann, F., 2015. Effects of the hydroedaphic gradient on tree species composition and aboveground wood biomass of oligotrophic forest ecosystems in the central Amazon basin. *Folia Geobot.* 50, 185–205. <http://dx.doi.org/10.1007/s12224-015-9225-9>.
- Thiers, B., 2016 [continuously updated]. Index Herbariorum: A global directory of public herbaria and associated staff. New York Botanical Garden's Virtual Herbarium. <<http://sweetgum.nybg.org/ih/>>.
- Toledo, J.J., Magnusson, W.E., Castilho, C.V., 2009. Influence of soil, topography and substrates on differences in wood decomposition between one-hectare plots in lowland tropical moist forest in Central Amazonia. *J. Trop. Ecol.* 25, 649–656. <http://dx.doi.org/10.1017/s0266467409990149>.
- UNFCCC, 1992. United Nations Framework Convention on Climate Change. In: United Nations Framework Convention on Climate Change (UNFCCC), Bonn, Germany. <<https://unfccc.int/resource/docs/convkp/conveng.pdf>>.
- UNFCCC, 1998. Kyoto Protocol to the United Nations Framework Convention on Climate Change. In: Document FCCC/CP/1997/7/Add.1 United Nations Framework Convention on Climate Change (UNFCCC), Bonn, Germany. <<http://unfccc.int/resource/docs/cop3/07a01.pdf>>.
- UNFCCC, 2015. Adoption of the Paris Agreement, Proposal by the President, Draft decision-CP.21600008831. In: United Nations Framework Convention on Climate Change (UNFCCC), Bonn, Germany. <http://unfccc.int/documentation/documents/advanced_search/items/6911.php?prirref=>>.
- Vale Jr., J.F., Nicodem, S., Melo, V.F., Uchôa, S.C.P., Cruz, D.L.S., 2016. Characterization of organic matter under different pedoenvironments in the Viruá National Park, in Northern Amazon. *Revista Brasileira de Ciência do Solo* 40, e0140480. <http://dx.doi.org/10.1590/18069657rbc20140480>.
- van Geffen, K.G., Poorter, L., Sass-Klaassen, U., van Logtstijn, R.S.P., Cornelissen, J.H. C., 2010. The trait contribution to wood decomposition rates of 15 Neotropical tree species. *Ecology* 91, 3686–3697. <http://dx.doi.org/10.1890/09-2224.1>.
- Vasconcelos, S.S., Fearnside, P.M., Graça, P.M.L.A., Nogueira, E.M., Oliveira, L.C., Figueiredo, E.O., 2013. Forest fires in southwestern Brazilian Amazonia: estimates of area and potential carbon emissions. *For. Ecol. Manage.* 291, 199–208. <http://dx.doi.org/10.1016/j.foreco.2012.11.044>.
- Wittmann, F., Schöngart, J., Parolin, P., Worbes, M., Piedade, M.T.F., Junk, W.J., 2006. Wood specific gravity of trees in Amazonian white-water forests in relation to flooding. *IAWA J.* 27, 255–268. <http://dx.doi.org/10.1163/22941932-90000153>.
- Yizhao, C., Jianyang, X., Zhengguo, S., Jianlong, L., Yiqi, L., Chengcheng, G., Zhaoli, W., 2015. The role of residence time in diagnostic models of global carbon storage capacity: model decomposition based on a traceable scheme. *Sci. Rep.* 5, 16155. <http://dx.doi.org/10.1038/srep16155>.
- Zanne, A., Lopez-Gonzalez, G., Coomes, D., Ilic, J., Jansen, S., Lewis, S., Miller, R., Swenson, N., Wiemann, M., Chave, J., 2009. Global wood density database. Dryad. Identifier: <<http://hdl.handle.net/10255/dryad.235>>.
- Zell, J., Kändler, G., Hanewinkel, M., 2009. Predicting constant decay rates of coarse woody debris: a meta-analysis approach with a mixed model. *Ecol. Model.* 220, 904–912. <http://dx.doi.org/10.1016/j.ecolmodel.2009.01.020>.